

Periodic Solutions of Certain Differential Equations with Quasibounded Nonlinearities

S. H. CHANG

*Department of Mathematics, The Cleveland State University, Cleveland,
Ohio 44115*

Submitted by W. F. Ames

1. INTRODUCTION

Consider the following nonlinear ordinary differential equation:

$$Lx + f(t, x, x', \dots, x^{(n-1)}) = p(t), \quad (1)$$

where L is an n th order linear differential operator with constant coefficients, f and p are continuous functions and periodic in t with period $T > 0$, and $\int_0^T p(u) du = 0$. The operator L is assumed to have the following decomposition:

$$L = D^m(D + a_1) \cdots (D + a_{n-m}), \quad (2)$$

where $D = d/dt$, $1 \leq m \leq n$, and for each $j = 1, \dots, n - m$, $a_j \neq i2\omega\pi/T$ for any integer ω . The function f in (1) is quasibounded [6]. We may define quasiboundedness as follows. Let us denote the norm of each $y = (y_1, \dots, y_n)$ in R^n by $\|y\| = \max_{j=1, \dots, n} |y_j|$. A function $f: R^{n+1} \rightarrow R$, $(t, y) \rightarrow f(t, y)$, which is T -periodic in t , is said to be *quasibounded* if the number

$$|f| = \min_{0 < \rho < \infty} \left(\max_{\substack{\|y\| \geq \rho \\ 0 \leq t < T}} \frac{|f(t, y)|}{\|y\|} \right) \quad (3)$$

is finite. When f is quasibounded, we call $|f|$ the *quasinorm* of f .

In this paper we shall prove the existence of T -periodic solutions for Eq. (1) by requiring f to be quasibounded and have a quasinorm smaller than certain positive number. This generalizes the results on existence of periodic solutions in Lazer [8] and Chang [2, 3]. Also, we shall prove a uniqueness theorem for the second order equation

$$x'' + cx' + f(t, x) = p(t), \quad (4)$$

which generalizes the uniqueness theorem of Leach [9, Theorem 2.1].

The proof of the existence theorem is based on Schauder's fixed point theorem [10] and on alternative methods (see e.g., Cesari [1], Hale [7]), and uses a technique generalizing those used in Lazer [8] and Chang [2, 3]. The proof of the uniqueness theorem for Eq. (4) is based on Sturm's comparison theorem. We shall state all results in Section 2 and prove them in Section 3.

2. THE RESULTS

Let X denote the Banach space of continuous T -periodic functions with the supremum norm, i.e. for each $x \in X$, $\|x\|_X = \max_{0 \leq t \leq T} |x(t)|$. For each $x \in X$ satisfying $\int_0^T x(u) du = 0$, define

$$A(x)(t) = \int_0^t x(u) du - \frac{1}{T} \int_0^T \int_0^s x(u) du ds. \quad (5)$$

It is easy to see that $A(x) \in X$, $\int_0^T A(x)(u) du = 0$, $A(x)'(t) = x(t)$, and $\|A(x)\|_X \leq T \|x\|_X$. Also, for each $x \in X$ satisfying $\int_0^T x(u) du = 0$, define

$$B(x)(t) = \int_0^t x(u) du. \quad (6)$$

Then $B(x) \in X$, $B(x)'(t) = x(t)$, and $\|B(x)\|_X \leq \frac{1}{2} T \|x\|_X$. For any positive integer k , let A^k denote the operator defined by $A^k(x) = A(\cdots A(x) \cdots)$, repeating k times.

Now, consider the operator L defined by (2). For any $x \in X$, define

$$H_j(x)(t) = (e^{a_j T} - 1)^{-1} \int_t^{T+t} e^{a_j(s-t)} x(s) ds, \quad (7)$$

where $j = 1, \dots, n - m$. It can be shown easily that $H_j(x) \in X$, $(D + a_j) H_j(x) = x$, and

$$\|H_j(x)\|_X \leq (1/\rho_j) \|x\|_X, \quad j = 1, \dots, n - m, \quad (8)$$

where $\rho_j = |\operatorname{Re}(a_j)|$ when $\operatorname{Re}(a_j) \neq 0$ and $\rho_j = (1/T) |e^{a_j T} - 1|$ when $\operatorname{Re}(a_j) = 0$ (note that $a_j \neq i2\omega\pi/T$).

Let S be the space of all x in X such that x is of class C^{n-1} . Then S becomes a Banach space under the norm

$$\|x\|_S = \max_{j=0,1,\dots,n-1} \|x^{(j)}\|_X, \quad \text{where} \quad x^{(j)} = d^j x / dt^j.$$

Then, for each $x \in X$ satisfying $\int_0^T x(u) du = 0$, $H_1 \cdots H_{n-m} B A^{m-1}(x) \in S$ and one can show that

$$\|H_1 \cdots H_{n-m} B A^{m-1}(x)\|_S \leq \sigma \|x\|_X, \quad (9)$$

where σ is a positive constant depending on $n, m, a_1, \dots, a_{n-m}, \rho_1, \dots, \rho_{n-m}$, and T . For example, when Eq. (1) is of the form

$$x^{(n)} + f(t, x, x', \dots, x^{(n-1)}) = p(t),$$

we consider, instead of the operator $H_1 \cdots H_{n-m} B A^{m-1}$, the operator $B A^{n-1}$ and find that $\|B A^{n-1}(x)\|_S \leq \sigma \|x\|_X$ with $\sigma = T^n/2$ when $T > 2$, $\sigma = T^{n-1}$ when $1 \leq T \leq 2$, and $\sigma = T$ when $0 < T < 1$. Also, when $f = f(t, x)$ in Eq. (1), we simply choose $S = X$ and find that (9) holds with $\sigma = T^m/(2\rho_1 \cdots \rho_{n-m})$.

Now, let us make the following assumptions:

(i) $f: R^{n+1} \rightarrow R$ is continuous, $p: R \rightarrow R$ is continuous, f and p are periodic in t with period $T > 0$, and $\int_0^T p(u) du = 0$.

(ii) There exists a number $M \geq 0$ such that for $x \in S$ one of the following two conditions holds:

(a) $\int_0^T f(s, x(s), x'(s), \dots, x^{(n-1)}(s)) ds \geq 0$ if $x(t) \geq M$ for all t , and $\int_0^T f(s, x(s), x'(s), \dots, x^{(n-1)}(s)) ds \leq 0$ if $x(t) \leq -M$ for all t .

(b) Integral in (a) ≤ 0 if $x(t) \geq M$ for all t , and ≥ 0 if $x(t) \leq -M$ for all t .

(iii) The function f is quasibounded as defined in (3) and has a quasinorm $|f| < \min\{\frac{1}{3}, 1/(6\sigma)\}$.

THEOREM 1. *If the assumptions (i), (ii), and (iii) are satisfied, then Eq. (1), with L given by (2), has at least one T -periodic solution.*

COROLLARY. *If we assume (i, ii), and*

$$(iv) \quad \lim_{\|y\| \rightarrow \infty} \frac{|f(t, y)|}{\|y\|} = 0 \text{ uniformly in } t,$$

then Eq. (1), with L given by (2), has at least one T -periodic solution.

Remark. Theorem 1 extends the result in [3, Theorem 1] concerning the equation $x^{(n)} + f(t, x) = p(t)$. The above corollary extends the results in Lazer [8] and [2] concerning second order cases.

Consider the second order equation

$$x'' + cx' + f(t, x) = p(t), \quad (4)$$

where c is any real number. Let us make the following assumptions:

(v) $f(t, x)$ and $(\partial/\partial x)f(t, x)$ are continuous for all (t, x) , $p(t)$ is continuous for all t , and $f(t, x)$ and $p(t)$ are T -periodic in t .

(vi) $f(t, x)$ satisfies one of the following conditions:

(a) $(\partial/\partial x)f(t, x) < 0$ for all (t, x) .

(b) $0 < (\partial/\partial x)f(t, x) < \frac{1}{4}c^2 + (2\pi/T)^2$ for all (t, x) .

(c) $\frac{1}{4}c^2 + (2n\pi/T)^2 < (\partial/\partial x)f(t, x) < \frac{1}{4}c^2 + (2(n+1)\pi/T)^2$ for all (t, x) and for some positive integer n .

THEOREM 2. *If the assumptions (v) and (vi) are satisfied, then Eq. (4) has at most one T -periodic solution.*

Remark. Theorem 2 extends the uniqueness theorem of Leach [9, Theorem 2.1] concerning the equation $x'' + f(x) = p(t)$.

3. THE PROOFS

Proof of Theorem 1. For $x \in S$, define

$$F(x)(t) = f(t, x(t), x'(t), \dots, x^{(n-1)}(t)) - \frac{1}{T} \int_0^T f(s, x(s), x'(s), \dots, x^{(n-1)}(s)) ds. \quad (10)$$

Then $F(x) \in X$ and $\int_0^T F(x)(u) du = 0$. Let R denote the set of real numbers. For any $\lambda_j \in R$ and $(x_j, r_j) \in S \times R$, $j = 1, 2$, let

$$\lambda_1(x_1, r_1) + \lambda_2(x_2, r_2) = (\lambda_1 x_1 + \lambda_2 x_2, \lambda_1 r_1 + \lambda_2 r_2).$$

Also, for each $(x, r) \in S \times R$, define $\|(x, r)\| = \|x\|_S + |r|$. Then $S \times R$ becomes a Banach space.

Suppose the condition (ii)-(a) holds. Define a mapping $P: S \times R \rightarrow S \times R$ by $P(x, r) = (\tilde{x}, \tilde{r})$ with

$$\tilde{x} = r + H_1 \cdots H_{n-m} B A^{m-1} (p - F(x)), \quad (11)$$

$$\tilde{r} = r - \frac{1}{T} \int_0^T f(s, \tilde{x}(s), \tilde{x}'(s), \dots, \tilde{x}^{(n-1)}(s)) ds. \quad (12)$$

It is clear that P is well-defined.

Since the quasinorm $|f| < \min\{\frac{1}{3}, 1/(6\sigma)\}$, there exists $\epsilon > 0$ such that $|f| + \epsilon < \min\{\frac{1}{3}, 1/(6\sigma)\}$. By the definition of quasiboundedness (3), there exists $\rho(\epsilon) > 0$ such that

$$|f(t, y)|/\|y\| < |f| + \epsilon \quad \text{whenever} \quad \|y\| \geq \rho(\epsilon) \quad \text{and} \quad 0 \leq t \leq T,$$

where $y = (y_1, \dots, y_n)$. Let

$$L = \max\{|f(t, y)| \mid 0 \leq t \leq T, \|y\| \leq \rho(\epsilon)\},$$

$$N = \max\left\{\frac{M}{1 - 3(|f| + \epsilon)}, \frac{M + 3\sigma\|p\|_X}{1 - 6\sigma(|f| + \epsilon)}, \frac{L}{|f| + \epsilon}, \rho(\epsilon)\right\},$$

and

$$C = \max\{(|f| + \epsilon)N, \sigma\|p\|_X + 2\sigma(|f| + \epsilon)N\}.$$

We note that $M + 3C \leq N$ and

$$|f(t, y)| \leq (|f| + \epsilon)N \quad \text{whenever} \quad \|y\| \leq N \quad \text{and} \quad 0 \leq t \leq T.$$

Now let

$$D = \{(x, r) \in S \times R \mid \|x\|_S \leq N, |r| \leq M + 2C\}.$$

Then D is a closed, bounded, and convex set in $S \times R$. It can be shown easily that $P(D) \subset D$, P continuous, and $P(D)$ relatively compact. Then by Schauder's fixed point theorem ([10], or see [5, p. 131]) there exists $(\theta, b) \in D$ such that $(\theta, b) = P(\theta, b) = (\tilde{\theta}, \tilde{b})$. From (11) and (12) we see that

$$\frac{1}{T} \int_0^T f(s, \theta(s), \theta'(s), \dots, \theta^{(n-1)}(s)) ds = 0$$

and hence by (10)

$$F(\theta)(t) = f(t, \theta(t), \theta'(t), \dots, \theta^{(n-1)}(t)).$$

Then it follows from $\theta - b = H_1 \cdots H_{n-m} B A^{m-1} (p - F(\theta))$ that

$$D^m(D + a_1) \cdots (D + a_{n-m}) \theta(t) + f(t, \theta(t), \theta'(t), \dots, \theta^{(n-1)}(t)) = p(t).$$

If the condition (ii)-(b) holds, we redefine the \tilde{r} in (12) as

$$\tilde{r} = r + \frac{1}{T} \int_0^T f(s, \tilde{x}(s), \tilde{x}'(s), \dots, \tilde{x}^{(n-1)}(s)) ds.$$

Then a similar argument as before establishes the desired result.

Proof of Corollary. The condition (iv) implies that $|f| = 0$.

Proof of Theorem 2. Suppose on the contrary that there exist two distinct T -periodic solutions $x_1(t)$ and $x_2(t)$ to Eq. (4). Then $\varphi(t) = x_2(t) - x_1(t)$ is a nontrivial T -periodic solution of

$$y'' + cy' + q(t)y = 0, \tag{13}$$

where

$$q(t) = \int_0^1 \frac{\partial}{\partial x} f[t, x_1(t) + s(x_2(t) - x_1(t))] ds.$$

Note that Eq. (13) is equivalent to

$$(e^{ct}y')' + e^{ct}q(t)y = 0. \quad (14)$$

Suppose the condition (vi)-(a) holds. Then $q(t) < 0$ and hence $e^{ct}q(t) < 0$. We claim that there exists t_0 such that $\varphi(t_0) = 0$ and $\varphi'(t_0) \neq 0$. For if $\varphi(t) > 0$ for all t , we have $(e^{ct}\varphi'(t))' = -e^{ct}q(t)\varphi(t) > 0$ and hence $e^{ct}\varphi'(t)$ strictly increasing, which is impossible for the nontrivial T -periodic solution $\varphi(t)$. Similarly it is impossible to have $\varphi(t) < 0$ for all t . Thus we have $\varphi(t_0) = \varphi(t_0 + T) = \dots = 0$. But, comparing Eq. (14) with the equation $(e^{ct}y')' = 0$, we obtain immediately a contradiction by Sturm's comparison Theorem (see e.g., [4, p. 208]).

Since the proof for the case (vi)-(b) is similar to both (vi)-(a) and (vi)-(c), we shall only prove (vi)-(c) in the following.

Suppose the condition (vi)-(c) holds. Then we have

$$e^{ct}[\frac{1}{4}c^2 + (2n\pi/T)^2] < e^{ct}q(t) < e^{ct}[\frac{1}{4}c^2 + (2(n+1)\pi/T)^2].$$

Comparing Eq. (14) with the equation

$$(e^{ct}y')' + e^{ct}[\frac{1}{4}c^2 + (2n\pi/T)^2]y = 0, \quad (15)$$

we conclude immediately that there exists t_0 such that $\varphi(t_0) = 0$ and $\varphi'(t_0) \neq 0$. Now, since Eq. (15) has a solution

$$y(t) = \exp(-\frac{1}{2}ct) \sin[2n\pi(t - t_0)/T]$$

with $2n+1$ zeros on $[t_0, t_0 + T]$, $\varphi(t)$ has at least $2n$ zeros in the open interval $(t_0, t_0 + T)$ by Sturm's comparison theorem. On the other hand, since the equation

$$(e^{ct}y')' + e^{ct}[\frac{1}{4}c^2 + (2(n+1)\pi/T)^2]y = 0$$

has a solution $y(t) = \exp(-\frac{1}{2}ct) \sin[2(n+1)\pi(t - t_0)/T]$ with $2n+1$ zeros in $(t_0, t_0 + T)$ and since $\varphi(t_0) = \varphi(t_0 + T) = 0$, we conclude that $\varphi(t)$ can have at most $2n$ zeros in $(t_0, t_0 + T)$. Hence, $\varphi(t)$ has exactly $2n$ zeros in $(t_0, t_0 + T)$. This implies that $\varphi'(t_0)$ and $\varphi'(t_0 + T)$ have opposite signs, a contradiction to that φ is a T -periodic solution. This completes the proof.

REFERENCES

1. L. CESARI, Functional analysis and periodic solutions of nonlinear differential equations, *Contributions to Differential Equations* 1 (1963), 149-187.
2. S. H. CHANG, Periodic solutions of certain second order nonlinear differential equations, *J. Math. Anal. Appl.* 49 (1975), 263-266.

3. S. H. CHANG, Periodic solutions of certain n -th order nonlinear differential equations, *Accad. Naz. Lincei Rend. Sci. Fis. Mat. Natur.* 5, **57** (1974), 519–524.
4. E. A. CODDINGTON AND N. LEVINSON, "Theory of Ordinary Differential Equations," McGraw-Hill, New York, 1955.
5. J. CRONIN, "Fixed Points and Topological Degree in Nonlinear Analysis," Math Survey 11, American Math. Society, Providence, R. I., 1964.
6. A. GRANAS, The theory of compact vector fields and some of its applications to topology of functional spaces (I), *Rozprawy Matematyczne* **30** (1962), 1–93.
7. J. K. HALE, "Ordinary Differential Equations," Wiley-Interscience, New York, 1969.
8. A. C. LAZER, On Schauder's fixed point theorem and forced second-order nonlinear oscillations, *J. Math. Anal. Appl.* **21** (1968), 421–425.
9. D. E. LEACH, On Poincaré's perturbation theorem and a theorem of W. S. Loud, *J. Differential Eqs.* **7** (1970), 34–53.
10. J. SCHAUDER, Der Fixpunktsatz in Funktionalraumen, *Studia Math.* **2** (1930), 171–180.